

SLAC-PUB-6670
CERN-TH.7451/94
TAUP-2201-94
OSU-RN-293/94
hep-ph/9409376

The Anomalous Magnetic Moments of the Electron and the Muon – Improved QED Predictions using Padé Approximants*

John Ellis[†], Marek Karliner[‡] and Mark A. Samuel[§]
Stanford Linear Accelerator Center
Stanford University, Stanford, California 94309, USA

and

Eric Steinfelds
Department of Physics
Oklahoma State University
Stillwater, Oklahoma 74078, USA

Abstract

We use Padé Approximants to obtain improved predictions for the anomalous magnetic moments of the electron and the muon. These are needed because of the very precise experimental values presently obtained for the electron, and soon to be obtained at BNL for the muon. The Padé prediction for the QED contribution to the anomalous magnetic moment of the muon differs significantly from the naive perturbative prediction.

*Work supported by the Department of Energy, contract DE-AC03-76SF00515.

[†]Permanent address: Theory Division, CERN, CH-1211, Geneva 23, Switzerland;
e-mail: johnne@cernvm.cern.ch

[‡]Permanent address: School of Physics and Astronomy, Raymond and Beverly Sackler Faculty of Exact Sciences, Tel Aviv University, Ramat Aviv, Tel Aviv, Israel; e-mail: marek@vm.tau.ac.il

[§]Permanent address: Department of Physics, Oklahoma State University, Stillwater, Oklahoma 74078, USA; e-mail: physmas@mvs.ucc.okstate.edu

Two of the most important tests of quantum electrodynamics (QED) are the comparisons between theory and experiment of the anomalous magnetic moments of the electron and the muon, a_e and a_μ respectively, where $a = (g-2)/2$. The latest Penning trap measurements of the electron and positron anomalies obtained by the University of Washington group¹ are:

$$a_{e-}^{\text{expt}} = 1159652188.4(4.3) \times 10^{-12} \quad (1)$$

and

$$a_{e+}^{\text{expt}} = 1159652187.9(4.3) \times 10^{-12} \quad (2)$$

The figures in brackets represent the error in the last 2 figures, a convention we will follow throughout this paper. Taking the average of eqs (1) and (2), one finds

$$a_e^{\text{expt}} = 1159652188.2(3.0) \times 10^{-12} \quad (3)$$

The most accurate measurement for the muon anomaly comes from the CERN $g-2$ experiment² in which it was found that

$$a_{\mu-}^{\text{expt}} = 1165936(12) \times 10^{-9} \quad (4)$$

and

$$a_{\mu+}^{\text{expt}} = 1165910(11) \times 10^{-9} \quad (5)$$

and the combined result is

$$a_\mu^{\text{expt}} = 1165923(9) \times 10^{-9} \quad (6)$$

where correlations are taken into account in combining the errors. A new $g-2$ muon experiment is being done at Brookhaven National Laboratory (BNL)³, and an improvement in the accuracy by a factor of about 20 is expected. In order to compare properly theory and experiment, one must improve correspondingly the accuracy of the theoretical predictions.

In an heroic feat, Kinoshita⁴ has calculated a_e in eighth order and Kinoshita, Nizic, Okamoto⁵ and Marciano⁶ have calculated a_μ in eighth order. Moreover, there have been some recent improvements in the analytic calculations^{7,8} of a_e and a_μ .

There have recently been several papers estimating coefficients in Perturbative Quantum Field Theory (PQFT) using Padé Approximants^{9,10}. This

procedure is known to give significant improvements on naive perturbative calculations in many condensed-matter applications¹¹, removes a large part of the discrepancy between experiment and QED calculations of the ortho-positronium decay rate^{10,12} and agrees well with other estimates of higher-order perturbative coefficients in QCD^{9,13}.

In this paper we will use Padé Approximants (PA's) to estimate, not just the next term in the perturbation series, but the entire sum of the series (as is frequently done in condensed-matter applications), for both a_e and a_μ . We obtain in this way a more accurate theoretical prediction of the QED contribution to a_μ , in particular, which lies outside the errors quoted previously.

The first step is to obtain an accurate value for the fine-structure constant α . The two most precise measurements of α are¹⁴

$$\alpha^{-1} = 137.0359979(32) \quad (7)$$

and¹⁵

$$\alpha^{-1} = 137.0359840(50) \quad (8)$$

We note that these two values differ by more than 2 standard deviations, but nevertheless take the average of eqs (7) and (8) to obtain

$$\alpha_{\text{exp}}^{-1} = 137.0359939(27) \quad (9)$$

The accuracy of this result limits the precision of tests of QED in the case of a_e , where both theory and experiment are extremely precise. The perturbation series for a_e is⁴

$$a_e = \frac{1}{2}\left(\frac{\alpha}{\pi}\right) - 0.328478965\left(\frac{\alpha}{\pi}\right)^2 + 1.17611(42)\left(\frac{\alpha}{\pi}\right)^3 - 1.434(138)\left(\frac{\alpha}{\pi}\right)^4 \quad (10)$$

and the error in the theoretical prediction is dominated by the error in α_{exp} .

The [N/M] Padé Approximant to a series

$$S = S_0 + S_1x + S_2x^2 + \dots + S_{N+M}x^{N+M} \quad (11)$$

is given by

$$[N/M] = \frac{a_0 + a_1x + \dots + a_Nx^N}{1 + b_1x + \dots + b_Mx^M} \quad (12)$$

where one chooses the coefficients a_i, b_j so that

$$[N/M] = S + O(x^{N+M+1}) \quad (13)$$

One can use such a PA either to predict the next coefficient S_{N+M+1} or to evaluate $[N/M]$ for the relevant value of x (in our case $x = \frac{\alpha}{\pi}$), and obtain an estimate for the sum of the series. Here we do the latter. The PA's are known to accelerate the convergence of many series by including the effects of higher (unknown) terms, thus providing a more accurate estimate of the series¹¹. The PA's also provide reliable estimates of many asymptotic series, as is the case in QED¹⁰ and QCD⁹.

For our application, we first construct PA's to a_e after removing an overall multiplicative factor of $(\frac{\alpha}{\pi})$. Our result for the $[1/2]$ PA is

$$[1/2] = 1159652169.1(24.0) \times 10^{-12} \quad (14)$$

and the $[2/1]$ PA agrees very well with the $[1/2]$:

$$[2/1] = 1159652169.0(24.0) \times 10^{-12} \quad (15)$$

The errors consist of 22.8 from α and 7.4 from the theoretical uncertainty. To obtain a_e one must add the contribution due to muon diagrams

$$\Delta a_e(\text{muon}) = 2.8 \times 10^{-12} \quad (16)$$

the contribution due to τ diagrams

$$\Delta a_e(\text{tau}) = 0.01 \times 10^{-12} \quad (17)$$

the hadronic contribution

$$\Delta a_e(\text{hadron}) = 1.6(2) \times 10^{-12} \quad (18)$$

and the purely weak contribution

$$\Delta a_e(\text{weak}) = 0.05 \times 10^{-12} \quad (19)$$

for a total of

$$\Delta a_e = 4.5(2) \times 10^{-12} \quad (20)$$

Thus the theoretical prediction for a_e is

$$a_e = 1159652173.5(24.0) \times 10^{-12} \quad (21)$$

Comparing eq (21) with eq (3), we see that there is beautiful agreement between theory and experiment:

$$a_e^{\text{expt}} - a_e = 14.7(24.0) \times 10^{-12} \quad (0.61\sigma) \quad (22)$$

As noted before, the error in eq (21) is dominated by the error in α . If one now assumes that QED is correct, and hence that theory and experiment agree, one obtains a new and more accurate value of α : α_{th} , where the [1/2] PA gives

$$\alpha_{\text{th}} = 137.03599228(86) \quad (23)$$

and the [2/1] PA leads to

$$\alpha_{\text{th}} = 137.03599227(86) \quad (24)$$

Comparing eq (23) with eq (9), one sees that there is beautiful agreement with the less-precise experimental value.

$$\alpha_{\text{expt}}^{-1} - \alpha_{\text{th}}^{-1} = 16(28) \times 10^{-7} \quad (0.57\sigma) \quad (25)$$

corresponding to the good agreement in eq (22). We note in passing that this provides an *a posteriori* justification for averaging naively the two most accurate measurements^{14,15} of α , and that the difference between the values of α_{th}^{-1} extracted using the perturbative series and the PA's is just 3×10^{-7} .

We now turn to the anomalous magnetic moment of the muon, a_μ . As is usual, we first consider the difference^{6,8}

$$a_\mu - a_e = 1.09433583(7)\left(\frac{\alpha}{\pi}\right)^2 + 22.869265(4)\left(\frac{\alpha}{\pi}\right)^3 + 127.00(41)\left(\frac{\alpha}{\pi}\right)^4 \quad (26)$$

In constructing a PA to this series, we must first remove an overall factor $(\frac{\alpha}{\pi})^2$ from the perturbative series. In this way, we obtain the [1/1] PA value

$$(a_\mu - a_e)[1/1] = 6194839(12) \times 10^{-12} \quad (27)$$

whereas the value from the series in eq (26) is

$$a_\mu - a_e = 6194791(12) \times 10^{-12} \quad (28)$$

Adding a_e from eq (21), after subtracting Δa_e from eq (20), we obtain

$$a_\mu^{\text{QED}} = 1165847008(12)(27) \times 10^{-12} \quad (29)$$

where the first error is due to numerical integrations used in evaluating the perturbative series, and the second error is due to α . This should be compared the value of Kinoshita and Marciano⁶

$$a_\mu^{\text{QED}} = 1165846955(44)(27) \times 10^{-12} \quad (30)$$

We note that the difference between these two estimates of a_μ^{QED} is considerably larger than the error propagated from α . The reason for our smaller error is that we have used the new more precise values in ref. [8].

If one now adds the hadronic¹⁶ and the weak⁶ contributions

$$\Delta a_\mu(\text{had}) = 7011(76) \times 10^{-11} \quad (31)$$

and

$$\Delta a_\mu(\text{weak}) = 195(10) \times 10^{-11} \quad (32)$$

one obtains the theoretical value

$$a_\mu = 116591907(77) \times 10^{-11} \quad (33)$$

The error is dominated by the error in $\Delta a_\mu(\text{had})$, and new, more precise experiments are underway in Novosibirsk and Frascati¹⁷ to reduce this error. Comparing eq (33) with eq (6), we obtain

$$a_\mu^{\text{expt}} - a_\mu = 4(9) \times 10^{-9} \quad (0.4\sigma) \quad (34)$$

The error in the difference between theory and experiment is dominated by the experimental error in eq (6), which should be reduced by a factor of 20 in the forthcoming BNL experiment³.

In summary, we have used PA to obtain new more precise values for the QED values of a_e and a_μ . These PA values, in effect, estimate the unknown higher-order contributions, and should be more precise than the naive perturbative values used previously. It would be interesting to compare our estimates with values obtained in a different way, for example using the effective charge approach¹³ which agrees very well with PA's in QCD applications⁹. Although smaller than some of the other uncertainties, the

shift we find in a_μ , in particular, is significantly larger than other theoretical uncertainties and the error due to α .

J. E., M. K. and M. S. would like to thank the theory group at SLAC for its very kind hospitality. The research of M.K. was supported in part by grant No. 90-00342 from the United States-Israel Binational Science Foundation (BSF), Jerusalem, Israel, and by the Basic Research Foundation administered by the Israel Academy of Sciences and Humanities. The research of M.A.S. was supported by the U.S. Department of Energy under Grant No. DE-FG02-94ER40852.

References

- [1] R. S. Van Dyck, Jr. P. B. Schwinberg and H. G. Dehmelt, Phys. Rev. Lett. 59, 26(1987).
- [2] J. Bailey et al , Phys. Lett. 68B, 191(1977).
- [3] Proposal for Brookhaven National Laboratory experiment E821, *A Precision Measurement of the Muon $g-2$ Value at the level of 0.35 ppm*, E. Benedict et al. (1986); co-spokesmen V.W. Hughes, W.M. Morse and B.L. Roberts.
- [4] T. Kinoshita, in Quantum Electrodynamics, edited by T. Kinoshita (World Scientific, Singapore, 1990), p.218.
- [5] T. Kinoshita, B. Nizic and Y. Okamoto, Phys. Rev. D41, 593(1990).
- [6] T. Kinoshita and W. J. Marciano, in Quantum Electrodynamics [3], p.419.
- [7] M. A. Samuel and G. Li, Phys. Rev. D44, 3935(1991) and D48, 1879(1993) erratum; G. Li, R. Mendel and M. A. Samuel, Phys. Rev. D47, 1723(1993).
- [8] S. Laporta and E. Remiddi, Phys. Lett. B301, 440(1993); S. Laporta and E. Remiddi, Phys. Lett. B265, 182(1991); S. Laporta, Phys. Lett. B312, 495(1993); S. Laporta, Nuovo Cimento 106, 675(1993); S. Laporta, Phys. Lett. B328, 522(1994); S. Laporta, Phys. Rev. D47, 4793(1993).

- [9] M. A. Samuel, G. Li and E. Steinfelds, Phys. Rev. D48, 869(1993); Phys. Lett. B323, 188(1994); M. A. Samuel, G. W. Li and E. Steinfelds, *On Estimating Perturbative Coefficients in Quantum Field Theory, Condensed Matter Theory and Statistical Physics*, Oklahoma State Univ. Research Note 278, August(1993); M. A. Samuel and G. W. Li, *Estimating Perturbative Coefficients in High Energy Physics and Condensed Matter Theory*, Oklahoma State Univ. Research Note 275(1992), to be published in Intl. Jrnl. of Theo. Phys.(1994).
- [10] M. A. Samuel and G. W. Li, Phys. Lett. B331, 114(1994).
- [11] For a review, see George A. Baker, Jr., *Essentials of Padé Approximants*, Academic Press (1975).
- [12] I.B. Khriplovich and A.I. Milstein, Budker Inst. report BINP-94-30, hep-ph/9404213.
- [13] A. L. Kataev and V. Starshenko, CERN-TH.7198/94, hep-ph/9405294 and CERN-TH.7400/94, hep-ph/9409395; see also P.M. Stevenson, Phys. Rev. D23, 2916(1981); G. Grunberg, Phys. Lett. 95B, 70(1980), E – *ibid.* 110B, 501(1982); Phys. Rev. D29, 2315(1984). For an alternative approach to reducing ambiguities in higher orders, see S.J. Brodsky, G.P. Lepage and P.B. Mackenzie, Phys. Rev. D28, 228(1983), S.J. Brodsky and H.J. Lu, SLAC-PUB-6481, hep-ph/9405218.
- [14] M. E. Cage et al, IEEE Trans. on Instrum. Meas. 38, 284(1989).
- [15] E. R. Williams et al, IEEE Trans on Instrum. Meas. 38, 233(1989).
- [16] L. Martinovic and S. Dubnicka, Phys. Rev. D42, 884(1990).
- [17] For a discussion of the feasibility of such an experiment, see V.W. Hughes and T. Kinoshita, Comments Nucl. Part. Phys. 14, 341(1985).